

hydrocarbons such as chloroform or methylene chloride; polar solvents such as tetrahydrofuran or N,N-dimethylformamide. Particularly, toluene and dichloromethane are preferred.

Page 59, please replace the paragraph beginning at line 20 with the following:

(Example 3) Production of 1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylsulfinyl-5-(1-oxy-pyridin-3-ylmethylamino)pyrazole-3-carbonitrile
(Compound No. 14)

Please replace the paragraph bridging pages 65 and 66 with the following:

In 350 ml and 110 ml of acetonitrile were dissolved 7.89 g (8.88 mmol) of bis(1-(2,6-dichloro-4-trifluoromethylphenyl)-3-carbonitrile-5-(pyrazin-2-ylmethylamino)pyrazol-4-yl)-disulfide, 5.44 g (31.6 mmol) of potassium trifluoromethanesulfinate, and 0.40 g (1.24 mmol) of dioxobis(acetylacetonato)molybdenum, and then 3.4 ml (27.2 mmol) of 80% t-butyl hydroperoxide solution was added dropwise thereto at room temperature. Furthermore, every 4 hours, the addition of 5.44 g (31.6 mmol) of potassium trifluoromethanesulfinate and 3.4 ml (27.2 mmol) of 80% t-butyl hydroperoxide solution was repeated twice, followed by stirring for 15 hours at room temperature. Then, 5.44 g (31.6 mmol) of potassium trifluoromethanesulfinate and 3.4 ml (27.2 mmol) of 80% t-butyl hydroperoxide solution were again added, followed by stirring for 6 hours. After removal of precipitated pale brown crystals by filtration, acetonitrile was removed from the filtrate by distillation under reduced pressure and ethyl acetate was added, followed by extraction. After washing of the organic layer with water, the solvent was removed under reduced pressure and the residue was subjected to a column chromatographic purification (Hex/AcOE₂ = 5/2) to obtain 0.88 g (1.7 mmol) of pale yellow crystals of 1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylsulfinyl-5-(pyrazin-2-ylmethylamino)pyrazole-3-carbonitrile in 9.6% yield.

Please replace the paragraph bridging pages 66 and 67 with the following:

ay
10028786-123004
Under a nitrogen atmosphere, 10.1 mg (0.24 mmol) of sodium borohydride was added to a methanol (1.5 ml) solution of 50 mg (0.11 mmol) of 1-(2,6-dichloro-4-trifluoromethylphenyl)-4-thiocyanato-5-(pyrazin-2-ylmethylamino)pyrazole-3-carbonitrile, followed by stirring at room temperature for 2 hours. After removal of the solvent from the reaction mixture by distillation under reduced pressure, 2 ml of DMF was added under nitrogen. After cooling in a dry ice-acetone bath, a DMF (0.5 ml) solution of 54.5 mg (0.11 mmol) of MEC-12, a trifluoromethylating agent manufactured by Daicel Chemical Industries, Ltd. was added thereto. After 1 hour of stirring at room temperature, ethyl acetate and water were added, followed by extraction. The organic layer was subjected to an LC analysis to observe the formation of 1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylsulfenyl-5-(pyrazin-2-ylmethylamino)pyrazole-3-carbonitrile in 41.7 area% yield.

Page 67, please replace the text at lines 20-23 with the following:

No. 7

as
1H NMR (CDCl₃): 4.66 (2H, d), 5.27 (1H, brs), 5.30 (1H, s), 5.61 (1H, s), 7.76 (2H, s), 8.41 (1H, d), 8.49 (1H, d), 8.54 (1H, s)

Page 68, please replace the text at lines 16-18 with the following:

No. 15

ab
1H NMR (CDCl₃): 4.84 (2H, d), 6.35 (1H, brs), 7.31 (3H, m), 7.77 (2H, s), 8.15 (1H, m)